

THE UNIVERSAL GENERATING FUNCTION OF ANALYTICAL POISSON STRUCTURES

BENOIT DHERIN

ABSTRACT. The notion of generating functions of Poisson structures was first studied in [2]. They are special functions which induce, on open subsets of $\mathbb{R}^d$, a Poisson structure together with the local symplectic groupoid integrating it. A universal generating function was provided in terms of a formal power series coming from Kontsevich star product. The present article proves that this universal generating function converges for analytical Poisson structures and compares the induced local symplectic groupoid with the phase space of Karasev–Maslov.

INTRODUCTION

Let U be an open subset of $\mathbb{R}^d$ and let S be a function defined on a neighborhood of $\{0\} \times \{0\} \times U$ in $(\mathbb{R}^d)^* \times (\mathbb{R}^d)^* \times U$. In [2] and in [5], it was shown that if S satisfies the SGS conditions and the SGA equation¹ (see Section 2), then S generates a Poisson structure on U given by

$$\alpha(x) := \frac{\partial^2 S}{\partial p_1 \partial p_2}(0, 0, x)$$

together with the local symplectic groupoid integrating it. Its structure maps, defined on a neighborhood $G_S(U)$ of U in T^*U , are given by

$$\begin{array}{lll} \epsilon(p, x) & = & (0, x) & \text{unit map} \\ i(p, x) & = & (-p, x) & \text{inverse map} \\ s(p, x) & = & \nabla_{p_2} S(p, 0, x) & \text{source map} \\ t(p, x) & = & \nabla_{p_1} S(0, p, x) & \text{target map.} \end{array}$$

Suppose now that we have a Poisson structure α on a manifold M and a local chart U with coordinates $x_1, \dots, x_d$. We may pose the reciprocal problem of finding a generating function on U such that $\alpha(x) := \frac{\partial^2 S}{\partial p_1 \partial p_2}(0, 0, x)$. Thus, solving this problem for α provides an explicit symplectic realization (in the sense of [11]) of the Poisson manifold (α, U) as well as an explicit description of the unique (up to isomorphism) local symplectic groupoid (see [12]) of (α, U) . As a consequence, observe that knowing the generating function S of α on U allows us to transform any Hamilton–Poisson system on U

$$\dot{X}(t) = \alpha(X(t)) \nabla f(X(t)), \quad X(0) = \xi \in U,$$

B. D. acknowledges partial support of SNF Grant No. 21-65213.01.

¹The SGA equation may be seen as the equation of an associative product in the operad of exact Lagrangian submanifolds of $(T^*\mathbb{R}^d)^k \times T^*\mathbb{R}^d$ where the compositions are given by symplectic reduction. See [3] for a presentation of this operad.

where $f \in C^\infty(U)$ into the following integrable Hamilton system on $G_S(U) \subset T^*U$

$$\begin{aligned}\dot{p}(t) &= -\nabla_x H_f(p(t), q(t)) \\ \dot{q}(t) &= \nabla_p H_f(p(t), q(t)),\end{aligned}$$

with Hamiltonian $H_f = -f \circ t$. The d constant of motions are given by the components of the source map $s^1, \dots, s^d$. In this case, results concerning integrable systems may be directly transfered to Hamilton–Poisson systems.

In fact, for any Poisson structure α and any local chart U , we may find a universal formal generating function for α . This is the main result of [2]. This universal generating function turns out to be the semi-classical part of Kontsevich star product associated to α on U ,

$$S_\epsilon(\alpha)(p_1, p_2, x) = (p_1 + p_2)x + \sum_{n=1}^{\infty} \frac{\epsilon^n}{n!} \sum_{\Gamma \in T_{n,2}} W_\Gamma \hat{B}_\Gamma(p_1, p_2, x).$$

This relates then symplectic groupoids with star products² as already foreseen by Coste, Dazord, and Weinstein in [4]. This universal formula is however not a “true” generating function in the sense that it is a formal power series in a formal parameter ϵ . The main result of this article is the following Theorem. It states that the universal generating function converges for analytical Poisson structures and compares the induced local symplectic groupoid with the construction of Maslov–Karasev [9].

Theorem 1. *Consider a Poisson manifold M with Poisson bivector α such that for any local chart U*

$$|\partial^\beta \alpha^{ij}(x)| \leq M_U(x)^{|\beta|+1},$$

where $\beta = (\beta_1, \dots, \beta_d) \in \mathbb{N}^d$ is a multi-index and $M_U(x)$ a positive function.

Then, the universal generating function

$$S_\epsilon(\alpha)(p_1, p_2, x) = (p_1 + p_2)x + \sum_{n=1}^{\infty} \frac{\epsilon^n}{n!} \sum_{\Gamma \in T_{n,2}} W_\Gamma \hat{B}_\Gamma(p_1, p_2, x),$$

converges absolutely for $\epsilon \in [0, 1)$ and

$$\|p_i\| \leq \frac{1}{64eM_U^2(x)}, \quad i = 1, 2.$$

Moreover, the local symplectic groupoid induced by S_ϵ on U is exactly the same – not only isomorphic – as the one constructed by Karasev and Maslov in [9]. In particular, if we consider the Poisson–Hamilton system

$$\dot{X}(t) = \alpha(X(t))p.$$

for small $p \in (\mathbb{R}^d)^*$, then the end-points of a solution X satisfy $X(0) = s(p, q)$ and $X(1) = t(p, q)$ for $q = \nabla_{p_2} S(-p, p, X(0)) = \nabla_{p_1} S(p, -p, X(1))$.

The proof of this Theorem is split between Propositions 1, 2 and 4. Note that the identification with Karasev local symplectic groupoid on U allows us to patch all the induced groupoids in the local charts into a global manifold $G(U)$, which is

²Since then, Karabegov in [6] has shown how to associate a formal symplectic groupoid to a general star product on a Poisson manifold. His approach differs essentially from ours by the fact that he defines a formal symplectic groupoid in terms of the functions on it.

the local groupoid integrating (M, α) . The question of defining a global generating function will be addressed elsewhere.

Organization of the article. In Section 1, we briefly review standard results on Poisson manifolds and local symplectic groupoids. Section 2 introduces the notion of generating functions of a Poisson structures and provides examples. In Section 3, we prove the convergence of the universal generating function for analytical Poisson structures. In Section 4, we compare the local symplectic groupoid generated by this convergent generating function and the one constructed by Maslov and Karasev in [9].

The results presented here are part of the author PhD thesis [5] and may be seen as a natural development of [2].

Acknowledgment. The author thanks Giovanni Felder and Alberto S. Cattaneo for their help and encouraging support as well as Alan Weinstein for useful suggestions.

1. LIE SYSTEM AND LOCAL SYMPLECTIC GROUPOIDS

In this section, we present a system of non-linear PDEs on an open subset U of $\mathbb{R}^d$ (that we called the reduced Lie system) and we briefly describe how a solution of this system generates a local symplectic groupoid structure over U . We also explain that any solution of the reduced Lie system may be used to transform a Poisson-Hamilton system on U into an integrable Hamilton system on the local symplectic groupoid. The results stated here were established by Coste, Dazord, Karasev, Maslov and Weinstein in [11], [4], [7], [8] and [9]. Therefore, we refer the reader to these articles for details and proofs.

The reduced Lie system. Let U be an open subset of $\mathbb{R}^d$ and V be an open neighborhood of U in T^*U . We say that the functions $s^i : V \rightarrow \mathbb{R}$, $i = 1, \dots, d$ are a solution of **the reduced Lie system** if for all $i, j = 1, \dots, d$

$$(1) \quad \sum_{u=1}^d \frac{\partial s^i}{\partial p^u} \frac{\partial s^j}{\partial x_u} - \frac{\partial s^i}{\partial x^u} \frac{\partial s^j}{\partial p_u} = \alpha^{ij}(s^1, \dots, s^d),$$

for some functions $\alpha^{ij} : U \rightarrow \mathbb{R}$, $i, j = 1, \dots, d$ and the $s = (s^1, \dots, s^d)$ satisfy the **initial condition** $s(0, x) = x$ and the **non-degeneracy condition**

$$\det(\nabla_x s(p, x)) \neq 0.$$

Remark 1. -

- (1) A necessary condition for the existence of a solution is that the matrix $\alpha(x) = (\alpha^{ij}(x))_{i,j=1}^d$ is a Poisson structure on U
- (2) If we denote by $\{\cdot, \cdot\}_\alpha$ the Poisson bracket on U associated to $(\alpha^{ij}(x))_{i,j=1}^d$ and by $\{\cdot, \cdot\}_\omega$ the Poisson bracket associated to the canonical symplectic form $\omega = dp \wedge dq$ on $V \subset T^*U$, then the reduced Lie system may be written as

$$\{s^i, s^j\}_\omega = \alpha^{ij}(s), \quad i, j = 1, \dots, d.$$

It implies that $s = (s^1, \dots, s^d)$ is a Poisson map from (V, ω) to (U, α) , i.e., that

$$\{s^*f, s^*g\}_\omega = s^*\{f, g\}_\alpha.$$

for all $f, g \in C^\infty(U)$. Such a map is also called a **symplectic realization** of the Poisson manifold (U, α) .

- (3) Suppose that s is a solution of the reduced Lie system. The initial and non-degeneracy conditions implies, by the implicit function theorem, that there exists an inverse Q such that

$$Q(p, s(p, q)) = q \quad \text{and} \quad s(p, Q(p, x)) = x.$$

Integrable Hamiltonian lift. Let α be a Poisson structure on $U \subset \mathbb{R}^d$ and let s be a solution of the reduced Lie system. An easy consequence of the fact that s is a Poisson map from (V, ω) to (U, α) is the following. The Hamilton-Poisson system on U with Hamiltonian $f \in C^\infty(U)$, i.e.,

$$\dot{X}(t) = \alpha^{ij}(X(t))\partial_j f(X(t)), \quad X(0) = \xi \in U,$$

may be lifted to the Hamilton system on V

$$\begin{aligned} \dot{p}(t) &= -\nabla_x H_f(p(t), q(t)) \\ \dot{q}(t) &= \nabla_p H_f(p(t), q(t)), \end{aligned}$$

with Hamiltonian $H_f = f \circ s$. In other words, if $(p(t), q(t))$ is the solution of the Hamilton system on V , then $X(t) := s(p(t), q(t))$ is the solution of the Poisson system on U provided that $s(p(0), q(0)) = \xi$. This Hamilton system is in fact an integrable one. Let us construct the d integrals of the motion as done in [8]. Consider on U the time-dependent Hamiltonian $K(x, t) = \langle Q(-pt, x), p \rangle$. Denote by ϕ_K^t the flow generated by K and define

$$t(p, q) := \phi_K^t(s(p, q))|_{t=1}.$$

In [8], it is shown that $t = (t^1, \dots, t^d)$ is the only map such that

$$\begin{aligned} (2) \quad \{t^i, t^j\}_\omega(p, x) &= -\alpha^{ij}(t(x, p)), \\ (3) \quad \{s^i, t^j\}_\omega(p, x) &= 0 \\ (4) \quad t(0, x) = x \quad \text{and} \quad \det(\nabla_x t(p, x)) &\neq 0. \end{aligned}$$

Thus, we may easily prove from (3) that $t = (t^1, \dots, t^d)$ constitute d constants of the motion of the Hamilton system on V . The system of PDEs defined by equations (1), (2) and (3) together with the respective initial and non-degeneracy conditions for s and t is called the **Lie system**.

Local groupoid structure. We may now describe the the local symplectic groupoid over U generated by a solution of the Lie system. Suppose that (s, t) is a solution of the Lie system defined on a open neighborhood $G_U(s, t)$ of U in T^*U . Thus, $G_U(s, t)$ inherits from T^*U the canonical symplectic form $dp \wedge dq$ turning it into a symplectic manifold. This will be the local symplectic groupoid over U associated to a solution of the Lie system. The structure maps are namely given by s for the **source**, by t for the **target** and by $\epsilon(x) = (0, x)$ for the **unit** $\epsilon : U \rightarrow G_U(s, t)$. The crucial point is to define the product and the inverse in the following way. Take $(p, x) \in G_U(s, t)$. If p is small enough, it is always possible to find a time-dependent Hamiltonian $f_\tau \in C^\infty(U)$, $\tau \in \mathbb{R}_+$, such that the flow Ψ_H^τ of its Hamiltonian lift $H = -f_\tau \circ t \in C^\infty(G_U(s, t))$ carries $(0, s(p, x))$ to (p, x) in time $\tau = 1$. Then, the **inverse** of (p, x) is defined as

$$(5) \quad i(p, x) = (\Psi_H^\tau)^{-1}|_{\tau=1}(0, t(p, x)),$$

and for any $(\bar{p}, \bar{x})$ such that $s(p, x) = t(\bar{p}, \bar{x})$, the **product** between the two points is defined as

$$(6) \quad m((p, x), (\bar{p}, \bar{x})) := \Psi_H^\tau((\bar{p}, \bar{x}))|_{\tau=1}.$$

This local symplectic groupoid is essentially unique, see for instance [4] and [8] for details and proofs.

2. GENERATING FUNCTIONS AND LOCAL SYMPLECTIC GROUPOIDS

In this section, we consider special functions which induce on an open subset U of $\mathbb{R}^d$ both a Poisson structure and the local groupoid which integrates it. As these functions are in fact usual generating functions of some Lagrangian submanifolds, we will call them generating functions of the local symplectic groupoid or of the associated Poisson structure. We will see how such generating functions provide explicit solutions of the Lie system. Let us first recall what we mean by generating functions in the context of cotangent bundle.

Definition 1 (Generating functions of exact Lagrangian submanifolds). *Consider the cotangent bundle T^*Q over a manifold Q endowed with its canonical symplectic form $\omega = -d\theta$. Then every 1-form $\mu \in \Omega^1(Q)$ can be considered as an embedding of Q in T^*Q ($\mu(Q) \subset T^*Q$). This embedding gives a Lagrangian submanifold when $\mu^*\omega = 0$, i.e. when μ is closed (namely $\mu^*\omega = -\mu^*d\theta = -d\mu^*\theta = -d\mu$). We can then associate to each closed 1-form a Lagrangian submanifold of T^*Q . Moreover when μ is exact $\mu = dS$, $S \in C^\infty(Q)$ we call S the generating function of $\mu(Q)$.*

We introduce some notations. Let $U \subset \mathbb{R}^d$ be an open set. Then set

$$B_n(U) := (\mathbb{R}^{d^*})^n \times U, \quad B_n^0(U) := \{0\}^n \times U \\ B_n := B_n(\mathbb{R}^d), \quad B_n^0 := B_n^0(\mathbb{R}^d).$$

Definition 2 (SGA and SGS). *Let $U \subset \mathbb{R}^d$ be an open subset. We say that a function $S \in C^\infty(B_2)$ satisfies the Symplectic Groupoid Associativity (SGA) equation on U if for each $(p_1, p_2, p_3, x) \in B_3(U)$ sufficiently close from $B_3^0(U)$ the following implicit equation for $\bar{x}, \bar{p}, \tilde{x}$ and $\tilde{p}$,*

$$\bar{x} = \nabla_{p_1} S(\bar{p}, p_3, x), \quad \bar{p} = \nabla_x S(p_1, p_2, \bar{x}), \\ \tilde{x} = \nabla_{p_2} S(p_1, \tilde{p}, x), \quad \tilde{p} = \nabla_x S(p_2, p_3, \tilde{x}),$$

has an unique solution and if the following additional equation holds

$$S(p_1, p_2, \bar{x}) + S(\bar{p}, p_3, x) - \bar{x}\bar{p} = S(p_2, p_3, \tilde{x}) + S(p_1, \tilde{p}, x) - \tilde{x}\tilde{p}.$$

Moreover, if $S(p, 0, x) = S(0, p, x) = px$ and $S(p, -p, x) = 0$, we say that S satisfies the Symplectic Groupoid Structure (SGS) conditions. At last, a function $S \in C^\infty(B_2)$ satisfying both the SGA equation and the SGS conditions will be called a symplectic groupoid generating function on U or simply, when no ambiguities arise, a generating function.

Let S be a generating function on U . We define the phase space of S on U by

$$G_U(S) := \left\{ (p, x) \in T^*U : \nabla_{p_1} S(0, p, x) \in U \quad \text{and} \quad \nabla_{p_2} S(p, 0, x) \in U \right\}.$$

It is easy to see from the SGS conditions that $G_U(S)$ is an neighborhood of the zero section in T^*U . Thus, $G_U(S)$ inherits the canonical symplectic structure of T^*U , turning $G_U(S)$ into a symplectic manifold. We proved in [2] and [5] that the matrix

$$\alpha(x) = \left(2\nabla_{p_k^1} \nabla_{p_l^2} S(0, 0, x) \right)_{k,l=1}^d$$

is a Poisson structure on U . This allows us to call S the **generating function of the Poisson structure** α . Moreover, the maps defined by

$$s(p, x) = \nabla_{p_2} S(p, 0, x) \quad \text{and} \quad t(p, x) = \nabla_{p_1} S(0, p, x)$$

constitute a solution of the Lie system on (U, α) . In particular s is a symplectic realization of the Poisson manifold (U, α) , i.e., a Poisson map from the symplectic manifold $(G_S(U), dp \wedge dq)$ into the Poisson manifold $(G_S(U), \alpha)$. The local inverses of s and t are given by the explicit formulas

$$Q(p, x) = \nabla_{p_2} S(-p, p, x) \quad \text{and} \quad \tilde{Q}(p, x) = \nabla_{p_1} S(p, -p, x).$$

Finally, summarizing the facts above, we have shown that $G_S(U)$ is the local symplectic groupoid integrating α with target t , source s , unit $\epsilon(x) = (0, x)$ and inverse $i(p, x) = (-p, x)$. The multiplication space $G_S(U)^{(m)}$ is the graph of dS .

Let us go through a series of examples.

Example 1. The function $S(p_1, p_2, x) = \langle x, \mu(p_1, p_2) \rangle$ where $\mu : \mathbb{R}^{d*} \times \mathbb{R}^{d*} \rightarrow \mathbb{R}^{d*}$ is a bilinear associative map on $\mathbb{R}^{d*}$ always satisfies the SGA equation. We get immediately that

$$\bar{x} = \langle x, \nabla_{p_1} \mu(\tilde{p}, p_2) \rangle, \quad \bar{p} = \mu(p_1, p_2), \quad \tilde{x} = \langle x, \nabla_{p_2} \mu(p_2, \tilde{p}) \rangle, \quad \tilde{p} = \mu(p_2, p_3).$$

An easy computation shows that that S satisfies the SGA equation iff

$$\mu(p_1, \mu(p_2, p_3)) = \mu(\mu(p_1, p_2), p_3).$$

The matrix

$$\nabla_{p^1} \nabla_{p^2} S(0, 0, x) = \left(2x^i \frac{\partial \mu_i}{\partial p_k^1 \partial p_l^2} (0, 0, x) \right)_{k,l=1}^d$$

is a Poisson structure if μ satisfies

$$\mu(p, 0) = \mu(0, p) = p \quad \text{and} \quad \mu(p, -p) = \mu(-p, p) = 0.$$

Notice that it is possible to find a function $S \in C^\infty(B_2)$ which satisfies the SGA equation but not the SGS conditions.

Example 2. Consider $T^*\mathbb{R}^2$ and identify $\mathbb{R}^{2*}$ with the complex plane. We define S as above with $\mu(p_1, p_2) = p_1 p_2$ induced by the complex multiplication. Thus,

$$S((p_x^1, p_y^1), (p_x^2, p_y^2), (x, y)) = x(p_x^1 p_x^2 - p_y^1 p_y^2) + y(p_x^1 p_y^2 + p_y^1 p_x^2)$$

satisfies the SGA equation but not the SGS conditions. By the way the matrix

$$\nabla_{p^1} \nabla_{p^2} S(0, 0, x) = \begin{pmatrix} x & y \\ y & -x \end{pmatrix}$$

is not a Poisson matrix.

Example 3. Let now be $\mu : \mathbb{R}^{d*} \times \mathbb{R}^{d*} \rightarrow \mathbb{R}^{d*}$ the vector addition. Then, this time, $S(p_1, p_2, x) = x(p_1 + p_2)$ satisfies not only the SGA equation but also the SGS conditions. The associate structure of symplectic groupoid is the trivial one. The structure maps are $s(p, x) = x$, $t(p, x) = x$, $\epsilon(x) = (0, x)$, $i(p, x) = (-p, x)$. The composition is the fiberwise addition. Moreover the induced Poisson structure on the base is the null Poisson structure.

Example 4. Consider $S(p_1, p_2, x) = x(p_1 + p_2) + \epsilon \alpha^{ij} p_i^1 p_j^2$ where $(\alpha^{ij})_{i,j=1}^d$ is a matrix. We get that

$$\bar{x}^l = x^l + \epsilon \alpha^{lj} p_j^3, \quad \bar{p}_l = p_l^1 + p_l^2, \quad \tilde{x}^l = x^l + \epsilon \alpha^{il} p_i^1, \quad \tilde{p}_l = p_l^2 + p_l^3.$$

One sees easily that S satisfies the SGA equation for any matrix $(\alpha^{ij})_{i,j=1}^d$. Moreover,

$$\nabla_{p^1} \nabla_{p^2} S(0, 0, x) = (\alpha^{ij})_{i,j=1}^d.$$

Imposing the SGS conditions to S is equivalent to say that the matrix $(\alpha^{ij})_{i,j=1}^d$ is skew-symmetric, which also imply that $\nabla_{p^1} \nabla_{p^2} S(0, 0, x)$ is a Poisson structure of constant rank. The multiplication space can then be described as

$$G_S^m = \left\{ \left((p_1, x + \epsilon \alpha p_2), (p_2, x - \epsilon \alpha p_1), (p_1 + p_2, x) \right), (p_1, p_2, x) \in B_2 \right\}.$$

The symplectic groupoid structure maps are given by

$$\begin{aligned} \epsilon(x) &= (0, x) \\ i(p, x) &= (-p, x) \\ s(p, x) &= x - \epsilon \alpha p \\ t(p, x) &= x + \epsilon \alpha p. \end{aligned}$$

Example 5. Suppose $\mathbb{R}^{d*}$ is given the structure of a Lie algebra, i.e, there is a Lie bracket $[\cdot, \cdot] : \mathbb{R}^{d*} \times \mathbb{R}^{d*} \rightarrow \mathbb{R}^{d*}$. In components we have $[p^1, p^2]_k = \alpha_k^{ij} p_i^1 p_j^2$ where the α_k^{ij} 's are the structure constants of the Lie algebra. The Baker–Campbell–Hausdorff formula

$$CBH(p_1, p_2) = p_1 + p_2 + \frac{1}{2}[p_1, p_2] + \frac{1}{12}([p_1, [p_2, p_2]] + [p_2, [p_2, p_1]]) + \dots,$$

induces a structure of group law in a neighborhood of 0 in $\mathbb{R}^{d*}$. As in Example 1, define

$$S_\epsilon(p_1, p_2, x) = \langle x, \frac{1}{\epsilon} CBH(\epsilon p_1, \epsilon p_2) \rangle.$$

S satisfies the SGA equation because of the associativity of CBH. Moreover, S satisfies also the SGS conditions. The induced Poisson structure on $\mathbb{R}^d$ is

$$\nabla_{p^1} \nabla_{p^2} S(0, 0, x) = (\alpha_k^{ij} x^k)_{i,j=1}^d.$$

Notice that, in this case, the induced Poisson structure is linear.

3. THE UNIVERSAL GENERATING FUNCTION

A generating function S produces a Poisson structure on U together with the local symplectic groupoid integrating it. Here, we investigate the reciprocal problem.

Question: Which Poisson structures α on U possess a generating function?

In [2], this problem was given a formal answer and a universal formal generating function was provided. It turns out to be the semi-classical part of Kontsevich star-product on U . In this section, we will first recall briefly this universal generating function. Then, we will prove that this universal generating function converges for analytical Poisson structure giving thus a true generating function, as well as an explicit solution of the Lie system.

Kontsevich, in [10], parametrizes all deformations (up to gauge equivalences) of the usual product of functions on an open set $U \subset \mathbb{R}^d$ in terms of the (formal) Poisson structures α (up to gauge equivalences) that one may put on U . He writes down the explicit parametrization

$$B_\epsilon(\alpha)(f, g)(x) = fg(x) + \sum_{n \geq 1} \frac{\epsilon^n}{n!} \sum_{\Gamma \in G_{n,2}} W_\Gamma B_\Gamma(\alpha)(f, g)(x),$$

where $G_{n,2}$ is a set of some special graphs, the Kontsevich graphs of type $(n, 2)$, the W_Γ are some real numbers associated to them and the $B_\Gamma(\alpha)$ are special bidifferential operators on $C^\infty(U)$.

The Proposition answering the previous question for an analytical Poisson structure is the following.

Proposition 1. *Consider a Poisson manifold M with Poisson bivector α such that for any local chart U*

$$|\partial^\beta \alpha^{ij}(x)| \leq M_U(x)^{|\beta|+1},$$

where $\beta = (\beta_1, \dots, \beta_d) \in \mathbb{N}^d$ is a multi-index and $M_U(x)$ a positive function.

Then, the universal generating function

$$S_\epsilon(\alpha)(p_1, p_2, x) = (p_1 + p_2)x + \sum_{n=1}^{\infty} \frac{\epsilon^n}{n!} \sum_{\Gamma \in T_{n,2}} W_\Gamma \hat{B}_\Gamma(p_1, p_2, x),$$

converges absolutely for $\epsilon \in [0, 1)$ and

$$\|p_i\| \leq \frac{1}{64eM_U^2(x)}, \quad i = 1, 2.$$

The reader may find definitions of Kontsevich graphs, weights and operators in the original paper [10] and a brief introduction to them as well as the definition of Kontsevich trees and symbols in [2]. Similar arguments may be found in [1] for the case of linear Poisson structures.

Proof of Proposition 1. This follows from a sequence of Lemmas.

Lemma 1. *Let $\Gamma \in T_{n,2}$, $p_1, p_2 \in B_\rho(0)$. Then,*

$$|\hat{B}_\Gamma(p_1, p_2, x)| \leq M^{2n-1} \rho^{n+1}.$$

Proof. The estimate follows from that $\hat{B}_\Gamma$ is a $(n+1)$ -homogeneous polynomial in the p variables and an n -homogeneous polynomial in α . Moreover, there is exactly $n-1$ derivatives differentiating the α s. $\square$

Lemma 2. $|T_{n,2}| \leq (16e)^n n!$

Proof. The graphs in $T_{n,2}$ are exactly the graphs in $G_{n,2}$ such that $\Delta(\Gamma)$ is a tree, i.e., a graph without cycle. Thus, a graph $\Gamma \in T_{n,2}$ may be described by specifying $\Delta(\Gamma)$, n^{n-2} choices, by giving each edge of $\Delta(\Gamma)$ an orientation, 2^{n-1} choices, by specifying for each vertex of $\Delta(\Gamma)$ if it has two outgoing edges going to $\{\bar{1}, \bar{2}\}$, just

one or none, 4^n choices, and, at last, by giving the labeling 1 or 2 for each edge, 2^n choices. This procedure of counting is by far not optimal. We count too many graphs and also graphs which are even not Kontsevich graphs. However, each graph in $T_{n,2}$ is counted one times. It gives us then the very crude estimate,

$$\begin{aligned} |T_{n,2}| &\leq n^{n-2} 2^{n-1} 4^n 2^n \\ &\leq n^n 16^n, \end{aligned}$$

which finishes the proof by remarking that $n^n \leq n!e^n$. $\square$

Definition 3. Let $\Gamma \in G_{n,2}$. An aerial vertex $v \in V_\Gamma^a$ is called a **terminal vertex** if it has no incoming edge. A terminal vertex $v \in V_\Gamma^a$ is called **terminal vertex of type 1** if one of $\gamma^1(v), \gamma^2(v)$ is in $\{\bar{1}, \bar{2}\}$ and the other is an aerial vertex. A terminal vertex $v \in V_\Gamma^a$ is called **terminal vertex of type 2** if both $\gamma^1(v)$ and $\gamma^2(v)$ are aerial vertices. Trivially, there can be no terminal vertex v such that $\gamma^1(v), \gamma^2(v) \in \{\bar{1}, \bar{2}\}$.

Lemma 3. Let be $\Gamma \in T_{n,2}$. Then, Γ has at least one terminal vertex.


Proof. Suppose Γ has no terminal vertex. Let us construct the following sequence of vertices. Take $v_0 \in V_\Gamma^a$. As v_0 is not a terminal vertex, we may chose for v_1 any vertex such that $(v_1, v_0) \in E_\Gamma$. As there is no terminal vertex, we can repeat this procedure infinitely many times. The result is an infinite sequence of vertices $\{v_l\}_{l \geq 0}$ of V_Γ^a . Now, as Γ is a Kontsevich tree, there must be no i, j such that $v_i = v_j$, this is a contradiction with the fact that V_Γ^a is a finite set. $\square$

Lemma 4. Let be $\Gamma \in T_{n,2}$. Then, we have the following alternative:


- (1) There is a terminal vertex of type 1, $v \in V_\Gamma^a$, such that $\Gamma_{|V_\Gamma \setminus \{v\}} \in T_{n-1,2}$.
- (2) There is a terminal vertex of type 2, $v \in V_\Gamma^a$, such that $\Gamma_{|V_\Gamma \setminus \{v\}} = \Gamma_k \Gamma_l$ where $\Gamma_k \in T_{k,2}$ and $\Gamma_l \in T_{l,2}$, $k + l = n - 1$.

Proof. Take $\Gamma \in T_{n,2}$. Consider v a terminal vertex in Γ . If v is of type 1 we are finished. If v is not of type 1, take another terminal vertex. If this other vertex is of type 1, we are finished. If there no terminal vertex of type 1, there must be at least one terminal vertex v of type 2. The restriction of $\Delta(\Gamma)$ to $V_\Gamma^a \setminus \{v\}$ separate $\Delta(\Gamma)$ in two disjoint components $\Delta_k(\Gamma)$ and $\Delta_l(\Gamma)$. They are disjoint otherwise there would be a cycle in $\Delta(\Gamma)$. Now, it is easy to see that Δ_k is a tree with k vertices and Δ_l a tree with l vertices such that $k + l = n - 1$ (if not $\Delta(\Gamma)$ is not a tree). Denote by Γ_k and Γ_l their associated Kontsevich trees. Then one has that $\Gamma_{|V_\Gamma \setminus \{v\}} = \Gamma_k \Gamma_l$. $\square$

We illustrate the two cases described by Lemma 4 by a picture.



This is best seen with a drawing:


$$\vec{\phi} := \psi_{\Gamma}(\vec{z}) = (\phi_1^1, \phi_1^2, \dots, \phi_n^1, \phi_n^2),$$

we have to show that there is at most 4^n configurations leading to the same $\vec{\phi}$. We do that by induction on n . For $n = 1$, clearly $|\psi_\Gamma^{-1}(\phi^1, \phi^2)| \leq 4$. Suppose that it is true for all Kontsevich trees in $T_{u,2}$ for $u = 1, \dots, n-1$. Then, Lemma 4, tells us that we have an alternative. If we are in the first case clearly by induction we have

$$|\phi_\Gamma^{-1}(\vec{\phi})| \leq 4^{n-1} \cdot 4 = 4^n$$

If we are in the second case, we have that,

$$|\phi_\Gamma^{-1}(\vec{\phi})| \leq 4^k 4^l 4 = 4^n.$$

□

Lemma 7. *Let $\Gamma \in T_{n,2}$ and W_Γ its Kontsevich's weight. Then,*

$$|W_\Gamma| \leq 4^n.$$

Proof. Consider the volume form

$$\omega_V = d\phi_1^1 \wedge d\phi_1^2 \wedge \dots \wedge d\phi_n^1 \wedge d\phi_n^2,$$

on $(S^1)^n$. Remark that

$$W_\Gamma = \frac{1}{(2\pi)^{2n}} \int_{\mathcal{H}^n \setminus D^n} \psi_\Gamma^* \omega_V.$$

Thus, by Lemma 6, we have that

$$|W_\Gamma| \leq 4^n \frac{V_{(S^1)^{2n}}}{(2\pi)^{2n}},$$

where $V_{(S^1)^{2n}} = (2\pi)^{2n}$ is the volume of $(S^1)^{2n}$. □

Now, supposing without restricting the generality, that $M_x > 1$, we can do the following estimate:

$$\begin{aligned} \left| \sum_{n=1}^{\infty} \epsilon^n \sum_{\Gamma \in T_{n,2}} W_\Gamma \hat{B}_\Gamma(p_1, p_2, x) \right| &\leq \sum_{n=1}^{\infty} \frac{\epsilon^n}{n!} (16e)^n n! 4^n M^{2n-1} \rho^{n+1} \\ &\leq \sum_{n=1}^{\infty} \epsilon^n (64eM^2)^n \rho^{n+1} \\ &\leq \frac{1}{64eM^2} \sum_{n=1}^{\infty} \epsilon^n, \quad p_1, p_2 \in B_\rho(0), \end{aligned}$$

which gives the desired result.

4. COMPARISON WITH KARASEV SYMMETRIC SOLUTION

We consider a Poisson manifold M with a Poisson structure α such that in every chart U for each $x \in U$ there exists a positive function $M_U(x)$ such that

$$|\partial^\beta \alpha^{ij}(x)| \leq M_U(x)^{|\beta|+1},$$

for all $i, j = 1, \dots, d$ and for all multi-index $\beta = (\beta_1, \dots, \beta_d) \in \mathbb{N}^d$.

We may assign to each chart U the generating function,

$$S_\epsilon(p_1, p_2, x) = x(p_1 + p_2) + \sum_{n=1}^{\infty} \frac{\epsilon^n}{n!} \sum_{\Gamma \in T_{n,2}} W_\Gamma \hat{B}_\Gamma(p_1, p_2, x)$$

which is convergent, thanks to Proposition 1, for (p_1, p_2, x) taken in a suitable neighborhood of $B_2^0(U)$.

In particular, there exists a neighborhood $G_U(S)$ such that $B_1^0(U) \subset G_U(S) \subset T^*U$ and on which

$$s(p, q) = \nabla_{p_2} S_\epsilon(p, 0, x) \quad \text{and} \quad t(p, q) = \nabla_{p_1} S_\epsilon(0, p, q),$$

take their values in U for all $(p, q) \in G_U(S)$.

For convenience, set $B(x) := 2\nabla_{p_1} \nabla_{p_2} S(0, 0, x) = 2\epsilon\alpha(x)$. The following Proposition tells us that the points $x_0 = s(p, q)$ and $x_1 = t(p, q)$ are the end points of a curve $x(t)$ in U satisfying a Hamilton-Poisson differential on U with the linear Hamiltonian $l_p(x) = px$ and initial condition $s(p, q)$.

Proposition 2. *Let be $(\bar{p}, \bar{q}) \in G_U(S)$. Consider the Poisson equation*

$$\dot{x}^i(t) = B^{ij}(x(t))\bar{p}_j, \quad x(0) = s(\bar{p}, \bar{q}).$$

Then,

$$x(1) = t(\bar{p}, \bar{q}).$$

*Moreover, the Hamilton system associated to $H(p, q) = -t^*l_p(p, q)$, $l_{\bar{p}}(x) = \langle \bar{p}, x \rangle$, on $G_U(S)$ takes the point $(0, s(\bar{p}, \bar{q}))$ to the point $(\bar{p}, \bar{q})$ in time $t = 1$.*

Proof. Consider the Hamilton function $l_{\bar{p}}(x) = \langle \bar{p}, x \rangle$ on U . The vector fields $-\omega^\sharp t^* dl_p$ and $B^\sharp dl_{\bar{p}}$ are t -related. Thus, the solution $(p(t), q(t))$ of the Hamilton system on $G_U(S)$,

$$(7) \quad \dot{p} = \left\langle \frac{\partial}{\partial q} t(p, q), \bar{p} \right\rangle, \quad p(0) = 0,$$

$$(8) \quad \dot{q} = -\left\langle \frac{\partial}{\partial p} t(p, q), \bar{p} \right\rangle, \quad q(0) = s(\bar{p}, \bar{q})$$

projects to the solution $x(t) = s(p(t), q(t))$ of the Poisson system $\dot{x} = B(x)p$, $x(0) = s(\bar{p}, \bar{q})$.

Remark now that for all q we have that

$$\begin{aligned} \left\langle \frac{\partial t}{\partial q}(t\bar{p}, q), \bar{p} \right\rangle &= \left\langle \nabla_x \nabla_{p_1} S_\epsilon(0, t\bar{p}, q), \bar{p} \right\rangle \\ &= \bar{p} + \sum_{i \geq 1} \epsilon^i \left\langle \nabla_x \nabla_{p_1} S_\epsilon^{(i)}(t\bar{p}, 0, q), \bar{p} \right\rangle \\ &= \bar{p}, \end{aligned}$$

as for $i \geq 1$, $S_\epsilon^{(i)}$ is a sum over Kontsevich's trees. Thus, if we insert $p(t) = \bar{p}t$ in (7) and in (8), Equation (7) is trivially always satisfied and remains for $q(t)$ the equation

$$\dot{q} = -\left\langle \frac{\partial t}{\partial p}(\bar{p}t, q), \bar{p} \right\rangle, \quad q(0) = s(\bar{p}, \bar{q}).$$

For small $\bar{p}$, there always exists a unique solution for $q(t)$ and as for all t we have that $s(p(t), q(t)) = s(\bar{p}, \bar{q})$, then

$$q(t) = Q(\bar{p}t, s(\bar{p}, \bar{q})).$$

Thus, $q(1) = Q(\bar{p}, s(\bar{p}, \bar{q})) = \bar{q}$ and we finally get that

$$t(\bar{p}, \bar{q}) = t(p(1), q(1)) = x(1)$$

□

In [7] and [8], Karasev gives another local solution of the Lie system on U . Take a point $x \in U$ and a point $p \in \mathbb{R}^{d*}$ small enough so that the Poisson system

$$\dot{x}^i = B^{ij}(x)p_j, \quad x(0) = x$$

has a solution $x(t)$ defined for $t \in [-1, 1]$. Consider the function

$$Q'(p, x) = \int_0^1 x(t) dt.$$

If p is small enough, we may inverse this function. We define the source of the point (p, q) as

$$s'(p, q) = x, \quad q = Q'(p, x).$$

The target is defined as $t'(p, x) = s'(-p, x)$.

It happens that (s', t') is a solution of the Lie system in a neighborhood of the null section. This solution is called **the symmetric solution**.

The next natural question to ask is how this symmetric solution compares to the one given by the generating function.

For that purpose, we will use a Proposition proved in [7] and [8] by Karasev. Before stating the Proposition, let us make a definition.

Definition 4. *Let (s, t) be a solution of the Lie system. Consider the following differential equation for $p(t)$,*

$$(9) \quad \dot{p} = \left\langle \frac{\partial t}{\partial q}(p, Q(p, x_0)), \bar{p} \right\rangle, \quad p(0) = 0, \quad x_0 \in U.$$

Then, define the exponential map associated to (s, t) as,

$$\exp_{x_0}(\bar{p}) := p(1).$$

Proposition 3 (Karasev [8]). *Let (s, t) and (s', t') be two solutions of the Lie system on U and let $\exp$ and $\exp'$ be their associated exponential maps. Then, the transformation*

$$\begin{aligned} p' &= \exp'_x(\exp_x^{-1})(p) \\ q' &= Q'(p', x), \quad x = s(p, q), \end{aligned}$$

is a symplectic map and satisfies

$$s'(p', q') = s(p, q), \quad t'(p', q') = t(p, q).$$

With the help of this Proposition, we may compare the symmetric solution and the one given by the generating function.

In fact, in [8], Karasev proves that for the symmetric solution (s', t') of the Lie system, $\exp'_x(p) = p$. As for the one given by the generating function, we get from the proof of Proposition 2 also that $\exp_x(p) = p$.

Proposition 4 (Comparison). *The the solution (s, t) of the Lie system given by the symplectic groupoid generating function is exactly the symmetric solution, i.e., if we consider the Poisson system*

$$\dot{x}^i(t) = B^{ij}(x)p_j, \quad x(0) = x,$$

then,

$$\begin{aligned} Q(p, x) &= \int_0^1 x(t) dt = \nabla_{p_2} S(-p, p, x), \\ Q(p, s(p, q)) &= q, \quad s(p, Q(p, x)) = x. \end{aligned}$$

Proof. Consider the symmetric solution (s', t') . In this case, by Proposition 3, the Karasev transformation relating (s, t) and (s', t') takes the form

$$\begin{aligned} p' &= p, \\ q' &= Q'(p, s(p, q)). \end{aligned}$$

This transformation is also symplectic. This imposes on the transformation that

$$\frac{\partial q'}{\partial q} = \text{id} \quad \text{and} \quad \frac{\partial q'}{\partial p} = \left(\frac{\partial q'}{\partial p} \right)^*.$$

Hence,

$$\frac{\partial q'}{\partial q} = \frac{\partial Q'}{\partial x}(p, x) \frac{\partial s}{\partial q}(p, q) = \text{id}.$$

As we also have that

$$\frac{\partial Q}{\partial x}(p, x) \frac{\partial s}{\partial q}(p, q) = \text{id},$$

then,

$$\frac{\partial Q'}{\partial x}(p, x) = \frac{\partial Q}{\partial x}(p, x),$$

for all p, x . Thus,

$$(10) \quad Q'(p, x) = Q(p, x) + C(p),$$

and the Karasev transformation becomes

$$\begin{aligned} p' &= p, \\ q' &= Q(p, s(p, q)) + C(p) = q + C(p). \end{aligned}$$

Now,

$$\frac{\partial q'}{\partial p} = \left(\frac{\partial q'}{\partial p} \right)^*,$$

implies that

$$\frac{\partial C}{\partial p}(p) = \left(\frac{\partial C}{\partial p}(p) \right)^*.$$

Hence, by Stokes, there exists a function $f : \mathbb{R}^{d*} \rightarrow \mathbb{R}$ such that $C(p) = \nabla f(p)$. Now, as we have for all p, x and $\lambda \in \mathbb{R}$ that,

$$\langle Q(\lambda p, x), p \rangle = \langle x, p \rangle, \quad \langle Q'(\lambda p, x), p \rangle = \langle x, p \rangle.$$

equation (10) gives that

$$\langle C(\lambda p), p \rangle = \langle \nabla f(\lambda p), p \rangle = 0.$$

As

$$\frac{d}{d\lambda} f(\lambda p) = \langle \nabla f(\lambda p), p \rangle = 0,$$

we get that $f(\lambda p) = f(0)$, i.e., f is constant and finally

$$\nabla f(p) = c(p) = 0$$

which completes the proof. $\square$

REFERENCES

- [1] Andler, M. ; Dvorsky, A. ; Sahi, S. Kontsevich quantization and invariant distributions on Lie groups. Ann. Sci. école Norm. Sup. (4) 35 (2002), no. 3, 371–390. **math.QA/9910104**.
- [2] Cattaneo, A. S.; Dherin, B. ; Felder, G. Formal symplectic groupoid. Comm. Math. Phys. 253 (2005), no. 3, 645–674., **math.SG/0312380**.
- [3] Cattaneo, A. S.; Dherin, B.; Felder, G. Formal Lagrangian operad. **math.SG/0505051**.
- [4] Coste, A. ; Dazord, P. ; Weinstein, A. Groupoides symplectiques. (French) [Symplectic groupoids] Publications du Département de Mathématiques. Nouvelle Série. A, Vol. 2, i–ii, 1–62, Publ. Dép. Math. Nouvelle Sér. A, 87-2, Univ. Claude-Bernard, Lyon, 1987.
- [5] Dherin, B. Star Products and Symplectic Groupoids. Diss. ETH Zürich. No. 15720 (2004). <http://www.math.ethz.ch/~dherin/diss.ps>
- [6] Karabegov, A. V. Formal symplectic groupoid of a deformation quantization. **math.QA/0408007** 41 pages. QA.
- [7] Karasev, M. V. Analogues of objects of the theory of Lie groups for nonlinear Poisson brackets. (Russian) Izv. Akad. Nauk SSSR Ser. Mat. 50 (1986), no. 3, 508–538, 638.
- [8] Karasev, M. V. The Maslov Quantization Conditions in Higher Cohomology and Analogs of Notions Developed in Lie Theory for Canonical Fibre Bundles of Symplectic Manifolds. II Selecta Mathematica Sovietica. Vol. 8 (1989), no. 3, 235–257.
- [9] Karasev, M. V.; Maslov, V. P. Nonlinear Poisson brackets. Geometry and quantization. Translated from the Russian by A. Sossinsky [A. B. Sosinskiĭ] and M. Shishkova. Translations of Mathematical Monographs, 119. American Mathematical Society, Providence, RI, 1993. xii+366 pp. ISBN: 0-8218-4596-9
- [10] Kontsevich, M. Deformation quantization of Poisson manifolds, I. **q-alg/9709040(1997)**.
- [11] Weinstein, A. The local structure of Poisson manifolds J. Diff. Geom. 18 (1983), 523–557.
- [12] Weinstein, A. Symplectic groupoids and Poisson manifolds. Bull. Amer. Math. Soc. (N.S.) 16 (1987), no. 1, 101–104.

D-MATH, ETH-ZENTRUM, CH-8092 ZÜRICH, SWITZERLAND

E-mail address: **dherin@math.ethz.ch**